

Behavioral Health 101: Schizophrenia

For Healthcare Providers

Role of PH Providers in Treating BH



Only 20% of adult clients with mental health disorders are seen by BH providers

Clients often prefer and receive treatment in primary care settings

60% of premature death in persons with Schizophrenia are due to medical conditions such as cardiovascular, pulmonary, and infection diseases

(American Hospital Association, 2014; Unützer, Harbin, Schoenbaum, & Druss, 2013)

Role of PCPs in Suicide Prevention



- 75% of individuals who die by suicide are in contact with a primary care physician in the year before their death
- 45% do so within one month of their death
- Only 20% of these patients saw a mental health professional in the preceding month
- 62% of antidepressant prescriptions in the U.S. are written by generalists (internists, pediatricians, PCPs)

(McDowell, Lineberry, & Bostwick, 2011; Office of the Surgeon General, US & National Action Alliance for Suicide Prevention, US. (2012)

Systematic integration facilitates the communication and coordination of:

- Physical healthcare
- Behavioral healthcare
- Substance use disorder treatment

Integrated care promotes a cohesive service delivery system & better continuity of care.

(SAHMSA, n.d.)

Common Disorders

- Depression
- Bipolar Disorders
- Anxiety Disorders
- Posttraumatic Stress Disorder
- Substance Use Disorders
- Schizophrenia & Psychotic Disorders

Schizophrenia & Psychotic Disorders



“They’ve always tried to help me, but having schizophrenia it was hard to receive the help.”

-Robert Lawton – film producer, writer, and director

“The problem is that you have a disease, but the disease is abnormal integrity, loyalty to a view of the world that the schizophrenic is willing to stake his life on.”

-Carl Whitaker – psychiatrist and therapist

Learning Objectives

- Recognize symptoms of schizophrenia and psychosis
- Identify 2 medications that are recommended for schizophrenia and psychosis
- List 2 treatment options for individuals who could have schizophrenia and psychosis
- Participate in and discuss an experiential exercise about schizophrenia

Schizophrenia

Prevalence

- 1.1% in the US population
- Higher prevalence in Medicaid population

Onset

- Age 16-30
- Primarily late adolescence
- Later for females than males

(Haffner et. al., 1994; National Institute for Mental Health, 2016)

Why to be aware

- Schizophrenia is a “serious mental illness” (SMI), which means it is chronic and has poor outcomes
- 50% to 90% of people with SMI have at least 1 chronic medical illness
- People with SMI die 25 years earlier than those without
- High need for integrated care

(Gold, Kilbourne, & Valenstein, 2008; Colton & Manderscheid, 2006)

“Natural Causes” of Death with SMI

- Heart disease
- Cancer
- Cerebrovascular, respiratory, and lung diseases



(Colton & Manderscheid, 2006)

Positive and Negative Symptoms

Symptoms of Schizophrenia: Positive and Negative

(Mental Health Guru Media, Inc., 2015)

Positive Symptoms: Psychosis

When a person has lost some contact with reality

Can occur

- Episodically, as a part of other conditions (Most common)
- Chronic in illnesses, such as schizophrenia
- In ranges (Mild to Severe)



Positive Symptoms: Psychosis

Delusions – fixed false beliefs

- Persecutory
- Grandiose
- Delusions of reference (e.g. TV talking to them)
- Thought insertion, control

Hallucinations – aberrant perceptions

- Visual
- Audio
- Tactile
- Olfactory

(Lane, 2015)

Positive Symptoms: Disordered Speech

- Switching topics rapidly
- Nonsensical speech (word salad)
- Making up words and sounds
- Comes from disordered thinking
 - Difficulty understanding information
 - Difficulty using information
 - Difficulty focusing
 - Problems with memory

(Mental Health America, 2015)



Negative Symptoms

- Social withdrawal
- Extreme apathy
- Lack of drive or initiative
- Emotional unresponsiveness



(Mental Health America, 2015)

Screening Instruments

None have been validated for use in primary care settings

- Low prevalence of the disorder
- Most screening questionnaires result in high false positive rates
- Prevention is still being researched

Screening Questions

- Use a normalizing lead in:
“Sometimes when people are [under stress/ feeling anxious/ feeling depressed], they can have strange experiences such as trouble with their thinking or seeing or hearing things that others don’t.”
- Follow up with “Tell me more” if patient answers “Yes”

(Viron, Bagget, Hill & Freudenreich, 2012)

Screening Questions

For delusions

- “Have you had any strange or odd experiences lately that are difficult to explain or that others would find hard to believe?”
- “Have you felt like people are watching or following you or that they want to harass or hurt you?”
- “Have you felt like others can hear your thoughts or that you can hear another person’s thoughts?”

(Viron et. al., 2012)

Screening Questions

For hallucinations

- “Have your eyes or ears ever played tricks on you?”
- “Have there been times when you heard or saw things that other people could not?”


(Viron et. al., 2012)

Treatment for Schizophrenia Disorders

- Treatment is ONLY for diminishing/eliminating symptoms
- Antipsychotic medication – primary treatment
- Psychosocial treatments
 - Illness management skills
 - Rehabilitation
 - Family education
 - Cognitive behavioral therapy
 - Self-help groups (e.g. NAMI Connections)


(NIMH, 2009)

Coordinated Specialty Care



(National Council for Behavioral Health, 2015)

Mental Health Treatment Team



(Viron et. al., 2012)

Medications: “Typical” Antipsychotics

- Chlorpromazine (Thorazine)
- Haloperidol (Haldol)
- Perphenazine (Etrafon, Trilafon)
- Fluphenazine (Prolixin)



Medications: Atypical Antipsychotics

- Risperidone (Risperdal)
- Olanzapine (Zyprexa)
- Quetiapine (Seroquel)
- Ziprasidone (Geodon)
- Aripiprazole (Abilify)
- Paliperidone (Invega)
- Asenapine (Saphris)
- Lurasidone (Latuda)

Group Activity



- Healthcare professional
 - Interviews patient with verbal screening questions
- Patient
 - Gets interviewed and answers “yes” to one of the questions
 - Answers other questions however they want to
- Observer
 - Evaluates interviewer
 - Times interaction
 - Gives feedback to interviewer
 - Asks interviewer what they would do
- Rotate roles
 - 6 minutes for each round

Process



- How difficult was it to conduct the screening?
- How long did it take to conduct the screening?
- What decisions did interviewers make after the screening?
- How could you use what you've learned for integrated patient care in your practice?

What Questions Do You Have?



Learning Objectives Revisited



- Recognize symptoms of schizophrenia and psychosis
- Identify 2 medications that are recommended for schizophrenia and psychosis
- List 2 treatment options for individuals who could have schizophrenia and psychosis
- Participate in and discuss an experiential exercise about schizophrenia

Reflection



- What are two or three of the important things you're taking away from this training?
- What changes will you make in your practice as a result of this training?

Parting Thought

“Never give up on someone with a mental illness. When ‘I’ is replaced by ‘We’, illness becomes wellness.”

Shannon L. Alder



References



- American Hospital Association (2014, February). Integrating behavioral health across the continuum of care. Chicago, IL: Health Research & Educational Trust. Retrieved from <http://www.hpoe.org/Reports-HPOE/Behavioral%20health%20FINAL.pdf>
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders 5th ed.). Washington, DC: Author.
- Anxiety and Depression Association of America (AADA). (2016). Find help: Treatment. Retrieved from <http://www.adaa.org/finding-help/treatment>
- Aldwin, C.M., Sutton, K.J., Chiara, G & Spiro III, A. (1996). Age differences in stress, coping, and appraisal: Findings from the Normative Aging Study. *Journal of Gerontology: Psychological Sciences*, 51(4), 179-188.
- Boyd, N. (n.d.). Symptoms of Schizophrenia: Positive and negative. Retrieved from <http://study.com/academy/lesson/symptoms-of-schizophrenia-positive-and-negative.html>
- Brady, K. T., Killeen, T. K., Brewerton, T., & Lucerini, S. (2000). Comorbidity of psychiatric disorders and posttraumatic stress disorder. *The Journal of Clinical of Psychiatry*, (suppl 7), 1-478.

References Continued...



- Brewin C.R, Andrews B., Valentine J.D. (2000). Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *J Consult Clin Psychol.* 68(5):748-66.
- Carlat, D. J. (1998). The psychiatric review of symptoms: a screening tool for family physicians. *American Family Physician,* 58(7), 1617-1624.
- Charney D.S. Psychobiological mechanisms of resilience and vulnerability: implications for successful adaptation to extreme stress. *Am J Psychiatry.* 2004 Feb; 161(2):195-216.
- Center for Quality Assessment and Improvement in Mental Health. (2007). STABLE resource toolkit. Retrieved from http://www.integration.samhsa.gov/images/res/STABLE_toolkit.pdf
- Colton, C. W., & Manderscheid, R. W. (2006). Congruencies in increased mortality rates, years of potential life lost, and causes of death among public mental health clients in eight states. *Prev Chronic Dis,* 3(2), A42.

References Continued...



- Earls, M. F. (2010). Incorporating recognition and management of perinatal and postpartum depression into pediatric practice. *Pediatrics*, 126(5), 1032-1039.
- Gold, K. J., Kilbourne, A. M., & Valenstein, M. (2008). Primary care of patients with serious mental illness: your chance to make a difference: a primary care visit may lead to regular care of side effects and comorbidities, especially if you coordinate care. *Journal of Family Practice*, 57(8), 515-526.
- Groberman, A. (2012). Difference between stress and anxiety. Retrieved on March 11, 2016 from <http://www.psyweb.com/articles/anxiety/difference-between-stress-and-anxiety>
- Hafner, H., Maurer, K., Loffler, W., & Fatkenheuer, B. an der Heiden W, Riecher-Rossler A, Behrens S, Gattaz WF. (1994). The epidemiology of early schizophrenia: Influence of age and gender on onset and early course. *Br. J. Psychiatry Suppl*, 23, 29-38.

References Continued...



- Hirschfeld, R. M. (2002). The Mood Disorder Questionnaire: a simple, patient-rated screening instrument for bipolar disorder. *Primary Care Companion J Clin Psychiatry*, 4(1), 9-11.
- Kessler R.C., Chiu W.T., Demler O., Walters E.E. (2005). Prevalence, severity, and comorbidity of twelve-month DSM-IV disorders in the National Comorbidity Survey Replication (NCS-R). *Archives of General Psychiatry*, 62(6):617-27.
- Kessler R.C., Berglund P.A., Demler O., Jin R., Walters E.E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication (NCS-R). *Archives of General Psychiatry*, 62(6):593-602.
- Kroenke, K., & Spitzer, R. L. (2002). The PHQ-9: a new depression diagnostic and severity measure. *Psychiatric annals*, 32(9), 509-515.
- Lane, C. (2015). Schizophrenia delusions. Retrieved from <http://www.schizophrenic.com/content/schizophrenia/symptoms/schizophrenia-delusions>

References Continued...



- Lang, A.J., Stein, M.B. (2005). An abbreviated PTSD checklist for use as a screening instrument in primary care. *Behaviour Research and Therapy*, 43, 585-594.
- Lang, A. J., Wilkins, K., Roy-Byrne, P. P., Golinelli, D., Chavira, D., Sherbourne, C., Rose, R. D., Bystritsky, A., Sullivan, G., Craske, M. G., & Stein, M. B. (2012). Abbreviated PTSD Checklist (PCL) as a Guide to Clinical Response. *General Hospital Psychiatry*, 34, 332-338.
- Lanius, R., Brand, B., Vermetten, E., & Spiegel, D. (2012). The dissociative subtype of posttraumatic stress disorder: Rationale, clinical and neurobiological evidence, and implementations. *Depression and Anxiety*, 29, 701-708. doi: 10.1002/da.21889.
- Mayo Clinic. (2010). How SSRI's work. Retrieved from <http://www.mayoclinic.org/diseases-conditions/depression/multimedia/antidepressants/vid-20084764>
- Mayo Clinic Staff. (2014). Generalized anxiety disorder: Lifestyle and home remedies. Retrieved from <http://www.mayoclinic.org/diseases-conditions/generalized-anxiety-disorder/basics/lifestyle-home-remedies/con-20024562>

References Continued...



- Mayo Clinic Staff. (2015). Diseases and conditions: Depression (major depressive disorder). Retrieved from <http://www.mayoclinic.org/diseases-conditions/depression/basics/definition/con-20032977>
- McDowell, A. K., Lineberry, T. W., & Bostwick, J. M. (2011). Practical Suicide-Risk Management for the Busy Primary Care Physician. *Mayo Clinic Proceedings*, 86(8), 792–800. <http://doi.org/10.4065/mcp.2011.0076>
- Mental Health America. (2015). Schizophrenia. Retrieved from <http://www.mentalhealthamerica.net/conditions/schizophrenia>
- Mental Health Guru Media, Inc. (2015). Symptoms of schizophrenia. Retrieved from <http://mental.healthguru.com/video/symptoms-of-schizophrenia>
- Merikangas, K. R., He, J. P., Burstein, M., Swanson, S. A., Avenevoli, S., Cui, L., & Swendsen, J. (2010). Lifetime prevalence of mental disorders in US adolescents: Results from the National Comorbidity Survey Replication–Adolescent Supplement (NCS-A). *Journal of the American Academy of Child & Adolescent Psychiatry*, 49(10), 980-989.

References Continued...



- National Council for Behavioral Health. (2015). First episode psychosis: Early intervention treatments for psychosis. Retrieved from <http://www.thenationalcouncil.org/topics/first-episode-psychosis/>
- National Institute of Mental Health. (2016). Anxiety Disorders. Retrieved from <http://www.nimh.nih.gov/health/topics/anxiety-disorders/index.shtml>
- National Institute of Mental Health. (2016). Schizophrenia. Retrieved from <http://www.nimh.nih.gov/health/statistics/prevalence/schizophrenia.shtml>
- National Institute for Mental Health. (2009). Schizophrenia. Retrieved from <http://www.nimh.nih.gov/health/publications/schizophrenia/index.shtml>
- Nestler Laboratory at the Icahn School of Medicine at Mount Sinai. (n.d.) Brain reward pathways. Retrieved from <http://neuroscience.mssm.edu/nestler/brainRewardpathways.html>

References Continued...



- Norman, S. B., Campbell-Sills, L., Hitchcock, C. A., Sullivan, S., Rochlin, A., Wilkins, K. C., & Stein, M. B. (2011). Psychometrics of a brief measure of anxiety to detect severity and impairment: The Overall Anxiety Severity and Impairment Scale (OASIS). *Journal of Psychiatric Research*, 45(2), 262–268. <http://doi.org/10.1016/j.jpsychires.2010.06.011>
- Norman, S. B., Hami Cissell, S., Means-Christensen, A. J. and Stein, M. B. (2006). Development and validation of an Overall Anxiety Severity And Impairment Scale (OASIS). *Depression and Anxiety*, 23, 245–249. doi: 10.1002/da.20182
- Norris, F.H. (1992). Epidemiology of trauma: Frequency and impact of different potentially traumatic events on different demographic groups. *Journal of Consulting and Clinical Psychology*, 60, 409-418.
- Office of the Surgeon General, US & National Action Alliance for Suicide Prevention, US. (2012). 2012 National Strategy for Suicide Prevention: Goals and Objectives for Action: A Report of the US Surgeon General and of the National Action Alliance for Suicide Prevention. Retrieved from http://www.surgeongeneral.gov/library/reports/national-strategy-suicide-prevention/full_report-rev.pdf

References Continued...



- Pfizer, Inc. (2002-2016). Patient health questionnaire (PHQ) screeners. Retrieved from <http://www.phqscreeners.com/>
- Prins, A. O., Ouimette, P., Kimerling, P., Cameron, R., Hugelshofer, R. P., Shaw-Hegwer, D. S., & Sheikh, J. J. (2003). The primary care PTSD screen (PC-PTSD): Development and operating characteristics. *Primary Care Psychiatry*, 9(1), 9-14.
- SAMHSA. (n.d.). What is integrated care?. Retrieved from <http://www.integration.samhsa.gov/about-us/what-is-integrated-care>
- SAMHSA. (n.d.). Medication assisted treatment. Retrieved from <http://www.samhsa.gov/medication-assisted-treatment>
- SAMHSA. (n.d.). Treatment for substance use disorders. Retrieved from <http://www.samhsa.gov/treatment/substance-use-disorders>
- Scheeringa, M.S., Zeanah, C.H., & Cohen, J.A. (2011). PTSD in children and adolescents: Toward an empirically based algorithm. *Depression and Anxiety*, 28, 770-782. doi:10.1002/da20736.

References Continued...



- Seitz, D. P. (2005). Screening mnemonic for generalized anxiety disorder. *Canadian Family Physician*, 51(10), 1340–1342.
- Skinner, H. A. (1982). The Drug Abuse Screening Test. *Addictive Behavior*, 7(4), 363–371.
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, 166(10), 1092-1097.
- University of Michigan Health System. (2011, August). Depression guideline update. Retrieved from <http://www.med.umich.edu/1info/FHP/practiceguides/depress/depress.pdf>
- Unützer, J., Harbin, H., Schoenbaum, M., & Druss, B. (2013). The collaborative care model: An approach for integrating physical and mental health care in Medicaid health homes. HEALTH HOME, Information Resource Center, 1-13. Retrieved from <https://www.medicaid.gov/State-Resource-Center/Medicaid-State-Technical-Assistance/Health-Homes-Technical-Assistance/Downloads/HH-IRC-Collaborative-5-13.pdf>

References Continued...



- U.S. Department of Veterans Affairs/Department of Defense. (2010). VA/DoD clinical practice guideline for management of post-traumatic stress. Retrieved from http://www.healthquality.va.gov/ptsd-sum_2010a.pdf (PDF)
- Viron, M., Baggett, T., Hill, M., & Freudenreich, O. (2012). Schizophrenia for primary care providers: how to contribute to the care of a vulnerable patient population. *The American Journal of Medicine*, 125(3), 223-230.
- Wisner, K. L., Parry, B. L., Piontek, C. M. (2002). Postpartum depression. *New England Journal of Medicine*, 347(3), 194-199.
- Zivin, K., Kim, H.M., McCarthy, J.F., Austin, K.L., Hoggatt, K.J., Walters, H.M., Valstein, M. (2007). Suicide mortality among individuals receiving treatment for depression in the VA health system: Associations with patient and treatment setting characteristics. *American Journal of Public Health*, 97(12), 2193-2198.

Resources

