

STANDARDIZED ONE PAGE PHARMACY PRIOR AUTHORIZATION FORM



Mississippi Division of Medicaid, Pharmacy Prior Authorization Unit,
550 High St., Suite 1000, Jackson, MS 39201

Magnolia Health/Envolve Pharmacy Solutions
Fax to: 1-866-399-0929 Ph: 1-866-399-0928
<https://www.magnoliahealthplan.com/providers/pharmacy.html>

UnitedHealthcare/OptumRx
Fax to: 1-866-940-7328 Ph: 1-800-310-6826
<http://www.uhccommunityplan.com/health-professionals/ms/pharmacy-program.html>

Medicaid Fee for Service/Change Healthcare
Fax to: 1-877-537-0720 Ph: 1-877-537-0722
<https://medicaid.ms.gov/providers/pharmacy/pharmacy-prior-authorization/>

BENEFICIARY INFORMATION

Beneficiary ID: _____ - _____ - _____ DOB: _____ / _____ / _____

Beneficiary Full Name: _____

PRESCRIBER INFORMATION

Prescriber's NPI: _____

Prescriber's Full Name: _____

Phone: _____

Prescriber's Address: _____

FAX: _____

PHARMACY INFORMATION

Pharmacy NPI: _____

Pharmacy Name: _____

Pharmacy Phone: _____

Pharmacy FAX: _____

CLINICAL INFORMATION

Requested PA Start Date: _____ Requested PA End Date: _____

Drug/Product Requested: _____ Strength: _____ Quantity: _____

Days Supply: _____ RX Refills: _____ Diagnosis or ICD-10 Code(s): _____

Hospital Discharge

Additional Medical Justification Attached

Medications received through coupons and/or samples are not acceptable as justification

PLEASE COMPLETE AND FAX DRUG SPECIFIC CRITERIA/ADDITIONAL DOCUMENTATION FORM FOUND BELOW

Prescribing provider's signature (signature and date stamps, or the signature of anyone other than the provider, are not acceptable)

I certify that all information provided is accurate and appropriately documented in the patient's medical chart.

Signature required: _____ Date: _____

Printed Name of Prescribing Provider: _____

FAX THIS PAGE

SUBMISSION AND/OR APPROVAL OF A DRUG PRIOR AUTHORIZATION REQUEST DOES NOT GUARANTEE MEDICAID PAYMENT FOR PHARMACY PRODUCTS OR THE AMOUNT OF PAYMENT. ELIGIBILITY FOR AND PAYMENT OF MEDICAID SERVICES ARE SUBJECT TO ALL TERMS AND CONDITIONS AND LIMITATIONS OF THE MEDICAID PROGRAM.

Confidentiality Notice: This communication, including any attachments, is for the sole use of the intended recipient(s) and may contain confidential and privileged information. Any unauthorized review, use, disclosure or distribution is prohibited. If you are not the intended recipient, please contact the sender by reply telephone (1-877-537-0722) or fax (1-877-537-0720) and destroy all copies of the original message. 05/05/2017

PRIOR AUTHORIZATION DESCRIPTION

Familial Hypercholesterolemia: REPATHA™ (evolocumab) and PRALUENT® (alirocumab)

Appendix A: Statin Contraindications

- Decompensated liver disease (symptoms can include jaundice, pruritus, ascites, variceal hemorrhage, or hepatic encephalopathy).
- Immune-mediated hypersensitivity to the HMG-CoA reductase inhibitor drug class (statins) as evidenced by an allergic reaction occurring with at least TWO different statins
- Laboratory-confirmed acute liver injury resulting from statin treatment
- Laboratory-confirmed rhabdomyolysis resulting from statin treatment
- Women who are breastfeeding, pregnant or are actively trying to become pregnant

Appendix B: Zetia Contraindications/Reasons to Discontinue

- Moderate or severe hepatic impairment (CP classes B and C)
- Women who are breastfeeding/pregnant or are actively trying to become pregnant
- Immune-mediated hypersensitivity to the cholesterol absorption as evidenced by an allergic reaction including anaphylaxis, angioedema, rash, or urticaria

Appendix C: A moderate-intensity statin may be more appropriate for the following adult populations if not able to tolerate a high-intensity statin

- Multiple or serious comorbidities, including impaired renal or hepatic function
- Unexplained ALT elevations >3 times ULN
- Active liver disease
- History of previous statin intolerance or statin-related muscle disorder
- Patient characteristics or concomitant use of drugs affecting statin metabolism
- ≥ 75 years of age
- History of hemorrhagic stroke
- Asian ancestry

Clinical atherosclerotic cardiovascular disease (ASCVD) includes:

- Acute coronary syndromes, or history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack, or peripheral arterial disease presumed to be of atherosclerotic origin.

PRIOR AUTHORIZATION DESCRIPTION

Low-Intensity Statin Therapy	Moderate-Intensity Statin Therapy	High-Intensity Statin Therapy
Daily dose lowers LDL-C by < 30%. on average	Daily dose lowers LDL-C by 30% to 50%. on average	Daily dose lowers LDL-C by ≥ 50%. on average
<ul style="list-style-type: none"> • Simvastatin 10 mg • Pravastatin 10-20 mg • Lovastatin 20 mg • Fluvastatin 20-40 mg • Pitavastatin (Livalo) 1 mg 	<ul style="list-style-type: none"> • Atorvastatin 10-20 mg • Rosuvastatin 5-10 mg • Simvastatin 20-40 mg • Pravastatin 40-80 mg • Lovastatin 40 mg • Fluvastatin XL (Lescol XL) 80 mg • Fluvastatin 40 mg twice daily • Pitavastatin (Livalo) 2-4 mg 	<ul style="list-style-type: none"> • Atorvastatin 40-80 mg • Rosuvastatin 20-40 mg

Criteria	Points
Family History	
First-degree relative with known premature* coronary and vascular disease, OR First-degree relative with known LDL-C level above the 95 th percentile	1
First-degree relative with tendinous xanthomata and/or arcus cornealis, OR Children aged < 18 years with LDL-C level above the 95 th percentile	2
Clinical History	
Patient with premature* coronary artery disease	2
Patient with premature* cerebral or peripheral vascular disease	1
Physical examination	
Tendinous xanthomata	6
Arcus cornealis prior to age 45 years	4
Cholesterol levels mg/dL (mmol/liter)	
LDL-C ≥330 mg/dL (≥8.5)	8
LDL-C 250 – 329 mg/dL (6.5 – 8.4)	5
LDL-C 190 – 249 mg/dL (5.0 – 6.4)	3
LDL-C 155 – 189 mg/dL (4.0 – 4.9)	1
DNA analysis	
Functional mutation in the <i>LDLR</i> , <i>apo B</i> or <i>PCSK9</i> gene	8
Diagnosis (diagnosis is based on total number of points obtained)	
Definite familial hypercholesterolemia	>8
Probable familial hypercholesterolemia	6 – 8
Possible familial hypercholesterolemia	3 – 5
Unlikely familial hypercholesterolemia	<3

*Premature – men < 55 years or women < 60 years Apo B= apolipoprotein B
 LDL-C= low density lipoprotein cholesterol; LDLR=low density lipoprotein receptor FH=familial hypercholesterolemia
 PCSK9=Proprotein convertase subtilisin/kexin type 9

CRITERIA/ADDITIONAL DOCUMENTATION

Homozygous Familial Hypercholesterolemia (HoFH)



BENEFICIARY INFORMATION					
Beneficiary ID: _____ - _____ - _____	DOB: ____/____/____				
Beneficiary Full Name: _____					
Homozygous Familial Hypercholesterolemia (HoFH) Criteria					
REPATHA™ (EVOLOCUMAB)					
Initial Approval Criteria for Repatha™ (evolocumab) may be approved when the following criteria are met:					
<input type="checkbox"/> Yes <input type="checkbox"/> No	The member is ≥ 13 years of age.				
AND					
<input type="checkbox"/> Yes <input type="checkbox"/> No	Evolocumab (Repatha) must be prescribed by or in consultation with a cardiologist, endocrinologist or lipid specialist and there is clinical documentation of one of the following: a.) Genetic confirmation of two mutant alleles at the LDLR, ApoB, PCSK9, or LDLRAP1 gene locus OR b.) Treated LDL-C of > 300 mg/dL or non-HDL-C 330 mg/dL or untreated LDL-C of > 500 mg/dL with either: i.) Cutaneous and/or tendon xanthoma before age 10 years OR ii.) Untreated LDL-C levels consistent with heterozygous familial hypercholesterolemia in both parents (> than 190 mg/dL);				
AND					
<input type="checkbox"/> Yes <input type="checkbox"/> No	Unable to meet LDL-C goal after treatment of at least 2 sequential 12-week trials of different high intensity statins [(i.e., atorvastatin ≥40mg or rosuvastatin ≥20mg] with at least one concomitant 12-week use of Zetia (ezetimibe) 10mg UNLESS contraindicated or not tolerated. (see Table 1 for dosages for therapy intensity; Appendix A for statin contraindications; Appendix B for Zetia contraindications) Adherence to the current statin regimen and Zetia (ezetimibe) must be evidenced by consistent pharmacy claims over the past 12 weeks, unless new to Medicaid. <ul style="list-style-type: none"> • If unable to tolerate a high-intensity statin, concomitant use with Zetia (ezetimibe) of a moderate to low-intensity statin at maximally tolerated dose can be used (see Appendix C). 				
AND					
<input type="checkbox"/> Yes <input type="checkbox"/> No	Repatha™ (evolocumab) will be used concomitantly with the statin and Zetia (ezetimibe), unless intolerance/contraindication justification is submitted (see Appendices A, B and C).				
<input type="checkbox"/> Yes <input type="checkbox"/> No	Repatha™ (evolocumab) is not being used concomitantly with Juxtapid® (lomitapide), Kynamro® (mipomersen), or another PCSK9 inhibitor.				
Recommended Dosing Regimen and Authorization Limit					
<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="text-align: left;">Drug</th> <th style="text-align: left;">Dosing Regimen</th> </tr> </thead> <tbody> <tr> <td>Repatha™(evolocumab)</td> <td>420mg SC once monthly</td> </tr> </tbody> </table>		Drug	Dosing Regimen	Repatha™(evolocumab)	420mg SC once monthly
Drug	Dosing Regimen				
Repatha™(evolocumab)	420mg SC once monthly				
Reauthorization Criteria:					
<input type="checkbox"/> Yes <input type="checkbox"/> No	Documentation of a LDL-C reduction from pretreatment level by ≥ 20% after adding Repatha™ for at least 90 days of therapy.				
AND					
<input type="checkbox"/> Yes <input type="checkbox"/> No	Is there clinical evidence of ongoing concomitant lipid lowering therapy (statin, ezetimibe, LDL-apheresis unless contraindicated/not tolerated)?				
Authorization					
Initial: If approved, initial coverage will be granted for up to 12 weeks.					
Maintenance: If approved, maintenance coverage will be reauthorized for periods of up to 52 weeks.					
APPENDICES AND TABLES CAN BE FOUND IN THE INSTRUCTION SHEET					

FAX THIS PAGE

SUBMISSION AND/OR APPROVAL OF A DRUG PRIOR AUTHORIZATION REQUEST DOES NOT GUARANTEE MEDICAID PAYMENT FOR PHARMACY PRODUCTS OR THE AMOUNT OF PAYMENT. ELIGIBILITY FOR AND PAYMENT OF MEDICAID SERVICES ARE SUBJECT TO ALL TERMS AND CONDITIONS AND LIMITATIONS OF THE MEDICAID PROGRAM.

Confidentiality Notice: This communication, including any attachments, is for the sole use of the intended recipient(s) and may contain confidential and privileged information. Any unauthorized review, use, disclosure or distribution is prohibited. If you are not the intended recipient, please contact the sender by reply telephone (1-877-537-0722) or fax (1-877-537-0720) and destroy all copies of the original message. 05/05/2017